
Recombinant human PEDF protein, His

Catalog Number: bs-41083P

Concentration: >0.5 mg/ml

AA Seq: 20-418/418

Predicted MW: 45.2

Detected MW: 55 kDa

Tags: His

Activity: Not tested

Endotoxin: Not analyzed

Purity: >95% as determined by SDS-PAGE

Purification: AC

Form: Lyophilized or Liquid

Storage: 20mM Tris-HCl (pH8.0) with 8M Urea.

Stored at -70°C or -20°C. Avoid repeated freeze/thaw cycles.

Background: Pigment epithelium derived factor, originally identified in conditioned medium of cultured human fetal retinal pigment epithelial (RPE) cells, is a neurotrophic protein that induces extensive neuronal differentiation in human Y79 retinoblastoma cells, a neoplastic counterpart of normal retinoblasts. It has been suggested that PEDF is synthesized by RPE cells and secreted into the retina interphotoreceptor matrix where it may influence development/differentiation of the neural retina. PEDF is a potent inhibitor of angiogenesis. As it does not undergo the S (stressed) to R (relaxed) conformational transition characteristic of active serpins, it exhibits no serine protease inhibitory activity. The PEDF gene is a member of the serpin gene family. Serpins are a group of serine protease inhibitors, some of which have also been reported to exhibit neurotrophic activity.

VALIDATION IMAGES



The purity of the protein is greater than 90% as determined by reducing SDS-PAGE.