

**bsm-33059M****[ Primary Antibody ]****NFKB p65 Mouse mAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**


|   |   |  |
|---|---|--|
| <b>Host:</b> Mouse<br><b>Clonality:</b> Monoclonal<br><b>GeneID:</b> 5970<br><b>Target:</b> NFKB p65<br><b>Purification:</b> affinity purified by Protein G<br><b>Concentration:</b> 1mg/ml<br><b>Storage:</b> Size : 50ul/100ul/200ul<br>0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.<br>Size : 200ug (PBS only)<br>0.01M PBS<br>Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.<br><b>Background:</b> NF-kappa-B is a ubiquitous transcription factor involved in several biological processes. It is held in the cytoplasm in an inactive state by specific inhibitors. Upon degradation of the inhibitor, NF-kappa-B moves to the nucleus and activates transcription of specific genes. NF-kappa-B is composed of NFKB1 or NFKB2 bound to either REL, RELA, or RELB. The most abundant form of NF-kappa-B is NFKB1 complexed with the product of this gene, RELA. Four transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2011]. | <b>Isotype:</b> IgG1<br><b>CloneNo.:</b> 12G3<br><b>SWISS:</b> Q04206 | <b>Applications:</b> <b>WB</b> (1:500-2000)<br><b>IHC-P</b> (1:200-1500)<br><b>IHC-F</b> (1:200-1500)<br><b>IF</b> (1:200-1500)<br><b>Reactivity:</b> Human, Mouse, Rat<br><br><b>Predicted MW.:</b> 61 kDa<br><b>Subcellular Location:</b> Cytoplasm ,Nucleus |
|---|---|--|

**— VALIDATION IMAGES —**

25 ug total protein per lane of various lysates (see on figure) probed with NFKB p65 monoclonal antibody, unconjugated (bsm-33059M) at 1:2000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Tonsil; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with NFKB p65 Monoclonal Antibody, Unconjugated (ascites of bsm-33059M) at 1:1500 overnight at 4°C, followed by conjugation to the bs-40296G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with NFKB p65 Monoclonal Antibody, Unconjugated (ascites of bsm-33059M) at 1:1500 overnight at 4°C, followed by conjugation to the bs-40296G-HRP and DAB (C-0010) staining.

**— SELECTED CITATIONS —**

- **[IF=6.7]** Yixu Wang. et al. Psoralen alleviates acute lung injury by covalently targeting Cys106 of HMGB1 in macrophages to inhibit inflammatory responses. PHYTOMEDICINE. 2025 Apr;;156807 WB ;Mouse. 40311596
- **[IF=4.868]** Zheng GH et al. Purple Sweet Potato Color Attenuates Kidney Damage by Blocking VEGFR2/ROS/NLRP3 Signaling in High-Fat Diet-Treated Mice. Oxid Med Cell Longev. 2019 Jan 22;2019:5189819. WB ;Mouse. 30805082
- **[IF=3.04]** Liu Z et al. The neuroprotective effect of lithium chloride on cognitive impairment through glycogen synthase kinase-3β inhibition in intracerebral hemorrhage rats.(2018) Eur J Pharmacol. 2018 Dec 5;840:50-59. WB ;Rat. 30336136