

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Desmoglein 2 Recombinant Rabbit mAb

Catalog Number: bsm-54310R
Target Protein: Desmoglein 2

Concentration: 1mg/ml

Form: Liquid Host: Rabbit

Clonality: Recombinant

Clone No.: 7D6
Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:50-200), IHC-F (1:50-200), IF (1:50-200)

Reactivity: Human
Predicted MW: 150 kDa
Entrez Gene: 1829
Swiss Prot: Q14126

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Store at -20°C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is

stable for at least two weeks at 2-4°C.

Background: Desmosomes are cell-cell junctions between epithelial, myocardial, and certain other cell

types. Desmoglein 2 is a calcium-binding transmembrane glycoprotein component of


desmosomes in vertebrate epithelial cells. Currently, three desmoglein subfamily members

have been identified and all are members of the cadherin cell adhesion molecule superfamily. These desmoglein gene family members are located in a cluster on

 $chromosome\ 18.\ This\ second\ family\ member\ is\ expressed\ in\ colon,\ colon\ carcinoma,\ and$

other simple and stratified epithelial-derived cell lines.

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Desmoglein 2 Monoclonal Antibody, Unconjugated(bsm-54310R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



(Negative control)Paraformaldehyde-fixed, paraffin embedded Human Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Desmoglein 2 Monoclonal Antibody, Unconjugated(bsm-54310R) at 1:200 overnight at 4° C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.