



Recombinant mouse CD160 protein, C-His Tag (HEK293)

Catalog Number: bs-43532P Concentration: >0.5 mg/ml

AA Seq: 28-160/185

Predicted MW: 19.9

Tags: C-His Tag
Activity: Not tested

Endotoxin: Not analyzed

Purity: >90% as determined by SDS-PAGE

Purification: AC

Form: Lyophilized or Liquid

Storage: PBS (pH7.4).

Stored at -70°C or -20°C. Avoid repeated freeze/thaw cycles.

Background: CD160 is an 27 kDa glycoprotein which was initially identified with the monoclonal antibody

BY55. Its expression is tightly associated with peripheral blood NK cells and CD8 T lymphocytes with cytolytic effector activity. The cDNA sequence of CD160 predicts a cysteine-rich, glycosylphosphatidylinositol-anchored protein of 181 amino acids with a single Ig-like domain weakly homologous to KIR2DL4 molecule. CD160 is expressed at the cell surface as a tightly disulfide-linked multimer. RNA blot analysis revealed CD160 mRNAs of 1.5 and 1.6 kb whose expression was highly restricted to circulating NK and T cells, spleen and small intestine. Within NK cells CD160 is expressed by CD56dimCD16+ cells whereas among circulating T cells its expression is mainly restricted to TCRgd bearing cells and to TCRab+CD8brightCD95+CD56+CD28-CD27-cells. In tissues, CD160 is expressed on all intestinal intraepithelial lymphocytes. CD160 shows a broad specificity for binding to both classical and nonclassical MHC class I molecules. [provided by RefSeq, Jul 2008]

VALIDATION IMAGES



The purity of the protein is greater than 55% as determined by reducing SDS-PAGE.