
FCAMR Rabbit pAb

Catalog Number: bs-16056R

Target Protein: FCAMR

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), ICC/IF (1:100-500), ELISA (1:5000-10000)

Reactivity: (predicted:Human)

Predicted MW: 55 kDa

Entrez Gene: 83953

Swiss Prot: Q8WWV6

Source: KLH conjugated synthetic peptide derived from human FCAMR: 1-100/532.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Fc (Ig constant fragment) receptors ensure protection of the host against foreign antigens, such as microorganisms and pathogens, by removing Ig-coated antigen complexes from circulation. Fc receptors are present on lymphoid and myeloid derivatives, where they mediate endocytosis of Ig-antigen complexes, antibody production in B cells through T cell antigen presentation, cytotoxicity and the release of cytokines and reactive oxygen species. The Fc α/μ receptor (FCAMR) functions as a receptor for the Fc fragment of IgA and IgM and mediates their endocytosis. FCAMR is a 532 amino acid protein that is expressed in mesangial cells and is upregulated by IFNG/IFN- γ . It is also expressed in secondary lymphoid organs, such as lymph node and appendix, kidney and intestine, suggesting an important role of the receptor for immunity in these organs.