bs-16063R

[Primary Antibody]

FCP1 Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 9150 SWISS: Q9Y5B0

Target: FCP1

Immunogen: KLH conjugated synthetic peptide derived from human FCP1:

201-300/961.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated


freeze/thaw cycles.

Background: This gene encodes a protein which interacts with the carboxy-

terminus of the RAP74 subunit of transcription initiation factor TFIIF, and functions as a phosphatase that processively dephosphorylates the C-terminus of POLR2A (a subunit of RNA polymerase II), making it available for initiation of gene expression. Mutations in this gene are associated with congenital cataracts, facial dysmorphism and neuropathy syndrome (CCFDN).

Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Feb 2011]

- VALIDATION IMAGES -



Sample: Heart(Mouse) Lysate at 40 ug Primary: Anti- FCP1 (bs-16063R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 104 kD Observed band size: 104 kD

Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FCP1) Polyclonal Antibody, Unconjugated (bs-16063R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

IHC-P (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500)

Reactivity: Mouse, Rat

(predicted: Human, Pig, Sheep, Cow, Chicken, Dog,

Horse)

Predicted

MW.: ¹⁰⁴ kDa

Subcellular Nucleus