bs-13995R

[Primary Antibody]

CORO1B Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET ––––––		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal	C C	IHC-P (1:100-500)
GenelD: 57175	SWISS: Q9BR76	IHC-F (1:100-500) IF (1:100-500)
Target: CORO1B		ICC/IF (1:100-500)
Immunogen: KLH conjugated synthetic peptide derived from human CORO1B: 301-400/489.		CORO1B: Reactivity: (predicted: Human, Mouse,
Purification: affinity purified by Protein A		Rat, Pig, Cow, Dog, Horse)
Concentration: 1mg/ml		
 Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Coronins are a family of WD repeat-containing, actin-binding proteins that localize to submembraneous areas and regulate cell motility and cytoskeletal rearrangement. Coronin 1A (COR01A, CLIPINA, CLABP, TACO, p57) can form coiled coil-mediated homotrimeric complexes that influence early phagosome formation. PKC-dependent phosphorylation of Coronin 1B (COR01B) at Serine 2 regulates leading edge dynamics and cell motility in fibroblasts through interactions with Arp2/3 complex. Coronin 1C (COR01C, Coronin 3, HCRNN4) is abundant in differentiating Neuro-2a cells, PC-12 cells and primary oligodendrocytes, where it is thought to influence neuron morphogenesis and migration. Coronin 2A (COR02A, CLIPINB, IR10, WDR2) is a component of the approximately 1.5-2 megadalton N-CoR (nuclear receptor corepressor) complex of 10-12 proteins, which recruits HDACs to generate repressive chromatin. Coronin 7 (COR07, CRN7) localizes to the Golgi membrane and influences the organization of intracellular membrane compartments and vesicular trafficking. Coronin 2B (COR02B, CLIPINC) and Coronin 6 (COR06) are similar to other members of this family, since they possess a conserved basic N-terminal motif and 3-10 WD repeats clustered in one to two core domains. 		MW.: ^{54 KDa} MW.: ^{54 KDa} Subcellular Cytoplasm Location: Cytoplasm PROIA, ed e 1B ind cell complex. n MIPINB, elex of sive lgi lar opin 2B o other basic N-

- SELECTED CITATIONS -----

• [IF=4.545] Ke Feng. et al. Compound Danshen Dripping Pill inhibits doxorubicin or isoproterenol-induced

cardiotoxicity. Biomed Pharmacother. 2021 Jun;138:111531 WB ;MOUSE. 10.1016/j.biopha.2021.111531