### bs-7478R

## [ Primary Antibody ]

# Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## PKN1 Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GenelD:** 5585 **SWISS:** Q16512

Target: PKN1

**Immunogen:** KLH conjugated synthetic peptide derived from human PKN1:

201-300/942.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene belongs to the protein kinase C

superfamily. This kinase is activated by Rho family of small G proteins and may mediate the Rho-dependent signaling pathway. This kinase can be activated by phospholipids and by limited proteolysis. The 3-phosphoinositide dependent protein kinase-1 (PDPK1/PDK1) is reported to phosphorylate this kinase, which may mediate insulin signals to the actin cytoskeleton. The proteolytic activation of this kinase by caspase-3 or related proteases during apoptosis suggests its role in signal transduction related to apoptosis. Alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul

2008]

Applications: IHC-P (1:100-500)

**IHC-F** (1:100-500) **IF** (1:100-500)

Reactivity: Mouse (predicted: Human,


Rat, Cow, Dog)

Predicted MW.: 104 kDa

Subcellular Cell membrane ,Cytoplasm

Location: , Nucleus

### VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PKN1) Polyclonal Antibody, Unconjugated (bs-7478R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

#### - SELECTED CITATIONS -

• [IF=1.785] Xia Wang. et al. Protein kinase N1 promotes proliferation and invasion of liver cancer. Exp Ther Med. 2021 Jun;21(6):1-9 IHC; Human. 33968181