
SLAP2 Rabbit pAb

Catalog Number: bs-13660R

Target Protein: SLAP2

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Mouse, Rat (predicted:Human, Rabbit, Sheep, Cow, Dog)

Predicted MW: 29 kDa

Entrez Gene: 84174

Swiss Prot: Q9H6Q3

Source: KLH conjugated synthetic peptide derived from human SLAP2: 101-200/261.

Purification: affinity purified by Protein A



Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: SLA2 is a 261 amino acid protein that exists as four alternatively spliced isoforms which localize to either the cytoplasm or to the cell membrane and contain one SH2 domain and one SH3 domain. Expressed predominately in tissues of the immune system, including thymus, spleen and lymph nodes, SLA2 functions as an adaptor protein that negatively regulates T-cell receptor (TCR) signaling and may inhibit T-cell activation. SLA2 interacts with Zap-70 and is subject to post-translational phosphorylation. The gene encoding SLA2 maps to human chromosome 20. Comprising approximately 2% of the human genome, chromosome 20 contains nearly 63 million bases that encode over 600 genes, some of which are associated with Creutzfeldt-Jakob disease, amyotrophic lateral sclerosis, spinal muscular atrophy, ring chromosome 20 epilepsy syndrome and Alagille syndrome.

VALIDATION IMAGES

Sample: Epithelium (Mouse) Lysate at 40 ug Primary: Anti- SLAP2 (bs-13660R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 29kD Observed band size: 29kD



Paraformaldehyde-fixed, paraffin embedded (rat spleen); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SLAP2) Polyclonal Antibody, Unconjugated (bs-13660R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining