

bs-11082R**[Primary Antibody]****Neurotrimin Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Pig, Sheep, Dog, Horse) Predicted MW.: 32 kDa Subcellular Location: Cell membrane
Clonality: Polyclonal		
GeneID: 50863	SWISS: Q9P121	
Target: Neurotrimin		
Immunogen: KLH conjugated synthetic peptide derived from human Neurotrimin/HNT: 101-200/344.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Cell adhesion molecules (CAMs) influence cell growth, differentiation, embryogenesis, immune response and cancer metastasis by networking information from the extracellular matrix to the cell. The four major families of cell adhesion molecules are immunoglobulin (Ig) superfamily (calcium-independent transmembrane glycoproteins), integrins (transmembrane non-covalently linked heterodimers of Alpha and Beta subunits), calcium-dependent cadherins and divalent cation-dependent selectins. Regulation of neuronal synaptic adhesion by CAMs has proven important for learning and memory. Proper embryonic morphogenic development is also heavily dependent on the regulation of cell adhesion molecules. Neurotrimin (hNT) is a neural cell adhesion molecule localizing to the cell membrane, where it acts as a lipid-anchor. Neurotrimin belongs to the IgLON family of proteins, a member of the larger immunoglobulin superfamily.		