

bs-4978R**[Primary Antibody]**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

PMCA1 Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: ELISA (1:5000-10000)
Clonality: Polyclonal		Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Pig, Cow, Dog, Horse)
GeneID: 490	SWISS: P20020	Predicted MW.: 138 kDa
Target: PMCA1		Subcellular Location: Cell membrane
Immunogen: KLH conjugated synthetic peptide derived from human PMCA1: 351-450/1258. < Extracellular >		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Plasma membrane-type Ca ²⁺ -ATPases (PMCA) mediate the export of bivalent calcium ions from eukaryotic cells. As members of the P class of ion-motive ATPases, PMCA are a functionally diverse group of proteins that are derived from alternatively spliced transcripts originating from four distinct genes, PMCA1, 2, 3, and 4. The expression of different PMCA isoforms and splice variants is regulated in a developmental, tissue- and cell type-specific manner, and with respect to the physiological needs of specific cell and tissue types. Spatial and temporal rates of resting intracellular Ca ²⁺ concentrations and Ca ²⁺ signaling in eukaryotic cells are dependent on the array of PMCA isoforms that are expressed in concert with the rate of Ca ²⁺ export. PMCA3 expression is confined to brain and skeletal muscle. The PMCA4 gene is located on human chromosome 1q25 and is ubiquitously expressed.		

— SELECTED CITATIONS —

- **[IF=4.01]** Balasubramaniam et al. Knockdown of sodium-calcium exchanger 1 induces epithelial-to-mesenchymal transition in kidney epithelial cells. (2017) J.Biol.Chem. 292:11388-11399 WB ;Dog. 28550085