bs-11619R

[Primary Antibody]

BIOSS ANTIBODIES www.bioss.com.cn sales@bioss.com.cn

RTP3 Rabbit pAb

- DATASHEET -

Clonality: Polyclonal GeneID: 83597 Target: RTP3

Immunogen: KLH conjugated synthetic peptide derived from human RTP3:

21-120/232.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Members of the RTP (receptor transporter proteins) family have

recently been discovered to influence bitter taste receptor expression in addition to inducing the expression of mammalian odorant receptors. RTP3 is a 232 amino acid single-pass type III membrane protein belonging to the TMEM7 family. Unlike other RTP proteins, RTP3 is not expressed in olfactory neurons but is expressed predominantly in liver. RTP3 is involved in the functional expression of bitter taste receptors and suppresses cell proliferation, and is also found in human circumvallate papillae and testis (regions where bitter taste receptors are expressed). The gene encoding RTP3 maps to human chromosome 3p21.31 within

C3CER1 (chromosome 3 common eliminated region 1), which is frequently eliminated in chromosomal deletions of solid tumors.

Host: Rabbit Isotype: IgG Ap

Applications: Flow-Cyt (0.2ug/test)

400-901-9800


Reactivity: Human (predicted: Mouse)

techsupport@bioss.com.cn

Predicted MW.: 27 kDa

Subcellular Location: Cell membrane

VALIDATION IMAGES



Blank control:Molt-4. Primary Antibody (green line): Rabbit Anti-RTP3 antibody (bs-11619R) Dilution: $1\mu g/10^{\circ}6$ cells; Isotype Control Antibody (orange line): Rabbit 1gG. Secondary Antibody: Goat anti-rabbit 1gG-PE Dilution: $1\mu g$ /test. Protocol The cells were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at at room temperature . Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.