bs-2084R

[Primary Antibody]

BIOSS ANTIBODIES www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn

Kir4.1 Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 3766 **SWISS:** P78508

Target: Kir4.1

Immunogen: KLH conjugated synthetic peptide derived from human Kir4.1:

81-180/379.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The KIR (for inwardly rectifying potassium channel) family of

potassium channels possess a greater tendency to allow potassium to flow into the cell rather than out of it. KIR4.1, also known as Kir1.2, is highly expressed in brain including glial cells, astrocytes and cortical neurons. KIR4.1 is also expressed in myelinsynthesizing oligodendrocytes and is crucial to myelination in the developing nervous system. The gene encoding human KIR4.1 maps to chromosome 1. KIR4.2, also known as Kir1.3, is expressed in kidney, lung, heart, thymus and thyroid during development. The gene encoding human KIR4.2 maps to chromosome 21 in the Down syndrome chromosome region 1, and KIR4.2 may play a role in the pathogenesis of Down's syndrome. KIR5.1 forms functional channels only by coexpression with either KIR4.1 or KIR4.2 in the kidney and pancreas. The gene encoding human KIR5.1 maps to

chromosome 17.

400-901-9800 **Applications: WB** (1:500-2000)

Reactivity: Mouse, Rat

(predicted: Human, Rabbit, Pig, Cow, Chicken, Dog,


Horse)

Predicted MW.: 42 kDa

1.144..

Subcellular Location: Cell membrane

VALIDATION IMAGES



25 ug total protein per lane of various lysates (see on figure) probed with Kir4.1 polyclonal antibody, unconjugated (bs-2084R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.

- SELECTED CITATIONS -

• [IF=3.77] Zhan Xie. et al. A novel model of subretinal edema induced by DL-alpha aminoadipic acid. EXP EYE RES. 2023 Jan;:109388 IF; Rabbit, Mouse, Rat. 36652968