

bs-6642R

[Primary Antibody]

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

FREAC3 Rabbit pAb


DATASHEET

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1 μ g/test) Reactivity: Human, Mouse, Rat (predicted: Cow, Chicken, Dog, Horse) Predicted MW.: 57 kDa Subcellular Location: Nucleus
Clonality: Polyclonal		
GeneID: 2296	SWISS: Q12948	
Target: FREAC3		
Immunogen: KLH conjugated synthetic peptide derived from human FOXC1/FREAC3: 101-200/553.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Binding of FREAC-3 and FREAC-4 to their cognate sites results in bending of the DNA at an angle of 80-90 degrees. Involvement in disease; Defects in FOXC1 are the cause of Axenfeld-Rieger syndrome type 3 (RIEG3); also known as Axenfeld-Rieger syndrome (ARS) or Axenfeld syndrome or Axenfeld anomaly. It is characterized by posterior corneal embryotoxon, prominent Schwalbe line and iris adhesion to the Schwalbe line. Other features may be hypertelorism (wide spacing of the eyes), hypoplasia of the malar bones, congenital absence of some teeth and mental retardation. When associated with tooth anomalies, the disorder is known as Rieger syndrome. Glaucoma is a progressive blinding condition that occurs in approximately half of patients with Axenfeld-Rieger malformations.		


VALIDATION IMAGES



Sample: Skin (Mouse) Lysate at 40 μ g Primary:
Anti-FREAC3 (bs-6642R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at
1/20000 dilution Predicted band size: 57 kD
Observed band size: 57 kD



Tissue/cell: rat brain tissue; 4%
Paraformaldehyde-fixed and paraffin-
embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block
endogenous peroxidase by 3% Hydrogen
peroxide for 30min; Blocking buffer (normal goat
serum, C-0005) at 37°C for 20 min; Incubation:
Anti-FOXC1/FREAC3 Polyclonal Antibody,
Unconjugated(bs-6642R) 1:200, overnight at 4°C,
followed by conjugation to the secondary
antibody(SP-0023) and DAB(C-0010) staining



Blank control:A431. Primary Antibody (green
line): Rabbit Anti-FREAC3 antibody (bs-6642R)
Dilution: 1 μ g /10⁶ cells; Isotype Control
Antibody (orange line): Rabbit IgG . Secondary
Antibody : Goat anti-rabbit IgG-AF647 Dilution:
1 μ g /test. Protocol The cells were fixed with 4%
PFA (10min at room temperature)and then
permeabilized with 90% ice-cold methanol for
20 min at -20°C. The cells were then incubated in
5%BSA to block non-specific protein-protein
interactions for 30 min at room temperature
.Cells stained with Primary Antibody for 30 min
at room temperature. The secondary antibody
used for 40 min at room temperature.
Acquisition of 20,000 events was performed.

SELECTED CITATIONS

- **[IF=2.766]** An et al. Chi-miR-4110 promotes granulosa cell apoptosis by targeting Sma- and Mad-related protein 2

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

(Smad2) in the caprine ovary. (2017) PLoS.One. 12:e0181162 WB ;Goat. 28704526