



Recombinant human Tau-4 protein, DsbC & His

Catalog Number: bs-41252P Concentration: >0.5 mg/ml

AA Seq: 46-211/352

Predicted MW: 45.2

Detected MW: 53 kDa

Tags: DsbC & His Activity: Not tested

Endotoxin: Not analyzed

Purity: >90% as determined by SDS-PAGE

Purification: AC

Form: Liquid

Storage: PBS (pH7.4) with 500mM NaCl (pH6.0).

Stored at -70°C or -20°C. Avoid repeated freeze/thaw cycles.

Background: Tau proteins are important Promotes microtubule assembly and stability, and might be

involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization. Tau proteins subcellular located in the axons of neurons, in the cytoso I and in association with plasma membrane components. It expressed in neurons. PNS-tau is expressed in the peripheral

nervous system while the others are expressed in the central nervous system.

VALIDATION IMAGES

kDa M R 130 — 95 — 70 — 53 — 40 — 33 — 25 —

The purity of the protein is greater than 90% as determined by reducing SDS-PAGE.