**bs-13256R** **[ Primary Antibody ]**

# GAB4 Rabbit pAb

www.bioss.com.cn
sales@bioss.com.cn
techsupport@bioss.com.cn
400-901-9800

## DATASHEET

**Host:** Rabbit **Isotype:** IgG

**Clonality:** Polyclonal

**GeneID:** 128954 **SWISS:** Q2WGN9

**Target:** GAB4

**Immunogen:** KLH conjugated synthetic peptide derived from human GAB4: 101-200/574.

**Purification:** affinity purified by Protein A

**Concentration:** 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** The Gab family of adaptor proteins function as molecular scaffolds that mediate protein recruit-ment to RTKs. Cytokine/growth factor triggering of protein tyrosine kinase receptors (RTKs) initiates signaling cascades that progress to the nucleus where signals for activation, proliferation and differentiation occur. This scaffolding mechanism represents a critical link in cytokine/growth factor signaling routes. Gab 1-4 contain Pleckstrin homology and potential binding sites for SH2 and SH3 domain-containing proteins. The recruitment of signaling partners to Gab family members is phosphorylation-dependent. Insulin receptor and EGF receptor signaling are among the cascades that rely on Gab family members to elicit a nuclear response to an extracellular stimulus. Gab 4 (GRB2-associated-binding protein 4), also designated GRB2-associated-binding protein 2-like (Gab 2-like), is a 574 amino acid protein that shares 62% sequence similarity with Gab 2 and contains one Pleckstrin homology domain.

**Applications:** **WB** (1:500-2000)
**IHC-P** (1:100-500)
**IHC-F** (1:100-500)
**IF** (1:100-500)
**ICC/IF** (1:100-500)
**ELISA** (1:5000-10000)

**Reactivity:** (predicted: Human, Mouse, Rat, Rabbit, Pig, Sheep, Cow, Dog, Horse)

**Predicted MW.:** 62 kDa

**Subcellular Location:** Cytoplasm