

bs-3526R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

CKMT2 Rabbit pAb**— DATASHEET —**

<p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>GeneID: 1160</p> <p>Target: CKMT2</p> <p>Immunogen: KLH conjugated synthetic peptide derived from human CKMT2: 331-419/419.</p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: CKMT2 (creatine kinase, mitochondrial 2 (sarcomeric)) is a creatine kinase isoenzyme, (reversibly) catalyzing the production of ATP from ADP by converting Phosphocreatine to creatine. Phosphocreatine serves as an energy reservoir for the rapid generation of ATP. Mitochondrial creatine kinase exists as two isoenzymes, sarcomeric mitochondrial creatine kinase (CKMT2) and ubiquitous creatine kinase; these are encoded by separate genes. Mitochondrial creatine kinases occurs in two different oligomeric forms: dimers and octamers, in contrast to the exclusively dimeric cytosolic creatine kinase isoenzymes. Sarcomeric mitochondrial creatine kinase has 80% homology with the coding exons of ubiquitous mitochondrial creatine kinase. The CKMT2 gene contains sequences homologous to several motifs that are shared among some nuclear genes encoding mitochondrial proteins and thus may be essential for the coordinated activation of these genes during mitochondrial biogenesis. Three transcript variants encoding the same protein have been found for this gene.</p>	<p>Isotype: IgG</p> <p>SWISS: P17540</p>	<p>Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)</p> <p>Reactivity: Rat (predicted: Human, Mouse, Rabbit, Pig, Cow, Dog, Horse)</p> <p>Predicted MW.: 46 kDa</p> <p>Subcellular Location: Cell membrane ,Cytoplasm</p>
---	--	---

— VALIDATION IMAGES —

Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-CKMT2 Polyclonal Antibody, Unconjugated(bs-3526R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat heart tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-CKMT2 Polyclonal Antibody, Unconjugated(bs-3526R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining