

## Recombinant Cyn. monkey CD3E protein, C-hFc (HEK293)

Catalog Number: bs-47075P

Concentration: >0.5 mg/ml

AA Seq: 22-117/198

Predicted MW: 36.9

Detected MW: Due to glycosylation, the protein migrates to 43-48 kDa based on Tris-Bis PAGE result.

Tags: C-hFc

Activity: Not tested

Endotoxin: <1.0 EU/μg as determined by LAL

Purity: >95% as determined by Tris-Bis PAGE

Purification: AC


Form: Lyophilized

Storage: Lyophilized from 0.22um filtered solution in PBS (pH7.4) with 5mM DTT. Normally 5% trehalose is added as protectant before Lyophilization.


Stored at -70°C or -20°C. Avoid repeated freeze/thaw cycles.

**Background:** CD3e molecule, epsilon is also known as CD3E, is a T-cell surface single-pass type I membrane glycoprotein. CD3E contains 1 Ig-like (immunoglobulin-like) domain and 1 ITAM domain. CD3E, together with CD3-gamma, CD3-delta and CD3-zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell development. CD3E plays an essential role in T-cell development, and defects in CD3E gene cause severe immunodeficiency. CD3E gene has also been linked to a susceptibility to type I diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.

### VALIDATION IMAGES



Cynomolgus CD3 Epsilon protein on Tris-Bis



Immobilized Cyno CD3 Epsilon, hFc Tag at

PAGE under reduced condition. The purity is greater than 95%.

1 $\mu$ g/ml (100 $\mu$ l/Well). Dose response curve for Biotinylated Anti-CD3E Ab. with the EC50 of 0.2 $\mu$ g/ml determined by ELISA.