
Recombinant human Notch 1 protein, C-His-Avi (HEK293)

Catalog Number: bs-47168P

Concentration: >0.5 mg/ml

AA Seq: 19-526/2555

Predicted MW: 56.5

Detected MW: Due to glycosylation, the protein migrates to 70-80 kDa based on Tris-Bis PAGE result.

Tags: C-His-Avi

Activity: Not tested

Endotoxin: <1.0 EU/μg as determined by LAL

Purity: >95% as determined by Tris-Bis PAGE; >95% as determined by SEC-HPLC

Purification: AC


Form: Lyophilized

Storage: Lyophilized from 0.22um filtered solution in PBS (pH7.4) with 5mM DTT. Normally 5% trehalose is added as protectant before Lyophilization.


Stored at -70°C or -20°C. Avoid repeated freeze/thaw cycles.

Background: This gene encodes a member of the Notch family. Members of this Type 1 transmembrane protein family share structural characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats, and an intracellular domain consisting of multiple, different domain types. Notch family members play a role in a variety of developmental processes by controlling cell fate decisions. The Notch signaling network is an evolutionarily conserved intercellular signaling pathway which regulates interactions between physically adjacent cells. In *Drosophila*, notch interaction with its cell-bound ligands (delta, serrate) establishes an intercellular signaling pathway that plays a key role in development. Homologues of the notch-ligands have also been identified in human, but precise interactions between these ligands and the human notch homologues remain to be determined. This protein is cleaved in the trans-Golgi network, and presented on the cell surface as a heterodimer. This protein functions as a receptor for membrane bound ligands, and may play multiple roles during development. [provided by RefSeq, Jul 2008].

VALIDATION IMAGES



Recombinant Human Notch 1 Protein on Tris-Bis PAGE under reduced conditions. The purity is greater than 95%.



The purity of Human Notch 1 Protein is greater than 95% as determined by SEC-HPLC.