

bs-6676R**[Primary Antibody]****GPR48 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 55366 Target: GPR48 Immunogen: KLH conjugated synthetic peptide derived from human LGR4: 65-170/951. < Extracellular > Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: G protein-coupled receptors (GPCRs), also designated seven transmembrane (7TM) receptors or heptahelical receptors, interact with G proteins (heterotrimeric GTPases) to synthesize intracellular second messengers, such as diacylglycerol, cyclic AMP, inositol phosphates and calcium ions. Their diverse biological functions range from vision and olfaction to neuronal and endocrine signaling and are involved in many pathological conditions. LGR4 (leucine-rich repeat-containing G protein-coupled receptor 4), also known as GPR48, is a 951 amino acid multi-pass membrane protein that contains 15 LRR (leucine-rich repeats) and belongs to the GPCR family. Expressed in multiple tissues, including testis, ovary, placenta, stomach, heart, kidney, pancreas and spleen, LGR4 functions as an orphan receptor that may be involved in physiologic activities throughout the cell. LGR4 is overexpressed in various cancer types and is thought to enhance carcinoma invasiveness and metastasis, suggesting an important role in tumor progression.	Isotype: IgG SWISS: Q9BXB1	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human (predicted: Mouse, Rat, Rabbit, Pig, Cow, Chicken, Dog, Horse) Predicted MW.: 104 kDa Subcellular Location: Cell membrane
--	---	--

— VALIDATION IMAGES —

Sample: HeLa Cell (Human) Lysate at 40 ug
 Primary: Anti-GPR48 (bs-6676R) at 1/300 dilution
 Secondary: HRP conjugated Goat-Anti-rabbit IgG (bs-0295G-HRP) at 1/5000 dilution Predicted band size: 104 kD Observed band size: 104 kD



Paraformaldehyde-fixed, paraffin embedded (Human cerebellum); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (GPR48 Rabbit pAb) Polyclonal Antibody, Unconjugated (bs-6676R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.