

bs-6674R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

KCNK3 Rabbit pAb**— DATASHEET —**

Host: Rabbit Clonality: Polyclonal GeneID: 3777 Target: KCNK3 Immunogen: KLH conjugated synthetic peptide derived from human KCNK3: 51-150/394. < Extracellular > Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: KCNK9 or TASK-3 (TWIK-related Acid sensitive K ⁺ channel) is a member of the potassium channel family of proteins that contain two-pore domain and four transmembrane domains. These channels are characterized as leak K ⁺ channels that are sensitive to changes in the extracellular pH. The physiological functions of TASK channels are largely unknown; it has been proposed that they may be involved in the regulation of breathing, aldosterone secretion and anesthetic-mediated neuronal activity. They were found to act in neurons' membrane potential and in resting K ⁺ currents. KCNK9 gene has been localized to the chromosomal region 8q24. The protein is expressed at high levels mainly in the brain and at low levels in other tissues. In contrast to normal tissues, it was found that KCNK9 is amplified in some human carcinomas such as breast, lung, colon and metastatic prostate. In 10% of breast cancer patients this gene is amplified, and in 44% the protein is over expressed. Monoclonal antibodies to KCNK9 are an important tool for studying the potassium channel family of proteins in different tissues.	Isotype: IgG SWISS: O14649 Applications: WB (1:500-2000) Reactivity: Mouse, Rat (predicted: Human, Rabbit, Sheep, Cow, Chicken, Dog) Predicted MW.: 43 kDa Subcellular Location: Cell membrane
---	--

— VALIDATION IMAGES —

Sample: Pancreas (Mouse) Lysate at 40 ug
Primary: Anti-KCNK3 (bs-6674R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 43 kD
Observed band size: 43 kD



Sample: Placenta (Mouse) Lysate at 40 ug
Primary: Anti-KCNK3 (bs-6674R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 43 kD
Observed band size: 43 kD

— SELECTED CITATIONS —

- **[IF=2.5]** Song Hai-chen. et al. TWIK-related acid-sensitive potassium channels TASK-1 and TASK-3 may participate in the process of the coexistence of asthma and OSA. SLEEP BREATH. 2023 Jul;;1-9 WB ;Mouse. 37428352