

**bs-6672R****[ Primary Antibody ]****GIRK1/KCNJ3 Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit <b>Clonality:</b> Polyclonal <b>GeneID:</b> 3760 <b>Target:</b> GIRK1/KCNJ3 <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human GIRK1: 81-180/501. <b>Purification:</b> affinity purified by Protein A <b>Concentration:</b> 1mg/ml <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. <b>Background:</b> This potassium channel is controlled by G proteins. Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium; as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium. This receptor plays a crucial role in regulating the heartbeat.	<b>Isotype:</b> IgG  <b>SWISS:</b> P48549	<b>Applications:</b> WB (1:500-2000)    <b>Reactivity:</b> Mouse, Rat (predicted: Human, Rabbit, Pig, Cow, Dog, GuineaPig)  <b>Predicted MW.:</b> 55 kDa  <b>Subcellular Location:</b> Cell membrane
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

**— VALIDATION IMAGES —**

Sample: Heart (Mouse) Lysate at 40 ug Primary: Anti-KCNJ3 (bs-6672R) at 1/2000 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 55 kD  
Observed band size: 60 kD



Sample: Heart (Mouse) Lysate at 40 ug Heart (Rat) Lysate at 40 ug Primary: Anti- GIRK1'KCNJ3 (bs-6672R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 55 kD Observed band size: 53 kD

**— SELECTED CITATIONS —**

- **[IF=5.9]** Zhao, Jing, et al. "Chronic obstructive sleep apnea causes atrial remodeling in canines: mechanisms and implications." Basic Research in Cardiology 109.5 (2014): 1-13. WB ;="Dog". 25015734