bs-0062R

[Primary Antibody]

CALB2 Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 794 **SWISS:** P22676

Target: CALB2

Immunogen: KLH conjugated synthetic peptide derived from human Calretinin:

211-271/271.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Calretinin is a calcium-binding protein which is abundant in auditory neurons. It belongs to the calbindin family. Calbindin 2 (calretinin), closely related to calbindin 1, is an intracellular calcium-binding protein belonging to the troponin C superfamily. Calbindin 1 is known to be involved in the vitamin-D-dependent calcium absorption through intestinal and renal epithelia, while the function of neuronal calbindin 1 and calbindin 2 is poorly understood. The sequence of the calbindin 2 cDNA reveals an open reading frame of 271 codons coding for a protein of 31,520 Da, and shares 58% identical residues with human calbindin1.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) **IF** (1:100-500) Flow-Cyt (2ug/Test)

Reactivity: Human, Mouse, Rat


(predicted: Pig, Cow, Dog,

Horse)


Predicted MW.: 29 kDa

Subcellular Location: Cell membrane ,Cytoplasm


VALIDATION IMAGES




Sample: Lane 1: Mouse Cerebrum Lysates Lane 2: Mouse Cerebellum Lysates Lane 3: Rat Cerebrum Lysates Primary: Anti- Calretinin (bs-0062R) at 1/1000 dilution Secondary: IRDve800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 29 kDa Observed band size: 29 kDa



Sample: U-87MG(Human) Cell Lysate at 30 ug Primary: Anti-Calretinin (bs-0062R) at 1/500 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 29 kD Observed band size: 29 kD



Tissue/cell: rat brain tissue: 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-Calretinin/CA Polyclonal Antibody, Unconjugated(bs-0062R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Blank control: A431, Primary Antibody (green line): Rabbit Anti-CALB2 antibody (bs-0062R) Dilution: 2µg/10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody: Goat anti-rabbit IgG-PE Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 0.1% PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

- [IF=3.9] Suzui, Masumi, et al. "Multiwalled carbon nanotubes intratracheally instilled into the rat lung induce development of pleural malignant mesothelioma and lung tumors." Cancer Science 107.7 (2016): 924-935. IHC; Rat. 27098557
- [IF=3.2] Li Xinming. et al. Platelet-rich fibrin promotes mesothelial cell proliferation and peritoneal repair by upregulating calretinin to prevent postoperative intestinal adhesion. INT J MED SCI. 2025 Feb;22(6):1254-1268 IF,WB;Mouse. 40084247