bs-0743R

[Primary Antibody]

BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn

DATASHEET -

Host: Rabbit **Isotype:** IgG

Clonality: Polyclonal

CTGF Rabbit pAb

SWISS: P29268

Target: CTGF

Immunogen: KLH conjugated synthetic peptide derived from mouse CTGF:

221-349/349.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene is a mitogen that is secreted by

vascular endothelial cells. The encoded protein plays a role in chondrocyte proliferation and differentiation, cell adhesion in many cell types, and is related to platelet-derived growth factor. Certain polymorphisms in this gene have been linked with a higher incidence of systemic sclerosis. [provided by RefSeq, Nov 2009].

400-901-9800 **Applications: WB** (1:500-2000)

Reactivity: Mouse, Rat

(predicted: Human, Rabbit,


Pig, Chicken, Dog)

Predicted MW.: 36 kDa

Subcellular Secreted ,Extracellular

Location: matrix

VALIDATION IMAGES



Sample: Lane 1: Mouse Kidney tissue lysates Lane 2: Rat Kidney tissue lysates Primary: Anti-CTGF (bs-0743R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 36 kDa Observed band size: 38 kDa

— SELECTED CITATIONS –

- [IF=8.46] Ciamporcero, E., et al. "YAP activation protects urothelial cell carcinoma from treatment-induced DNA damage." Oncogene (2015). WB ;Human. 26119935
- [IF=6.854] Ciamporcero et al. YAP activation protects urothelial cell carcinoma from treatment-induced DNA damage. (2016) Oncogene. 35:1541-53 WB; Human. 26119935
- [IF=4.6] Jiaxu Zhou. et al. JTE-013 Alleviates Pulmonary Fibrosis by Affecting the RhoA/YAP Pathway and Mitochondrial Fusion/Fission. PHARMACEUTICALS-BASE. 2023 Oct;16(10):1444 IHC,IF,WB;Mouse. 37895915
- [IF=5.3] Mya Thandar. et al. Mesenchymal stem cells derived from adipose tissue and umbilical cord reveal comparable efficacy upon radiation-induced colorectal fibrosis in rats. AM J CANCER RES. 2024 Apr 15;14(4):1594-1608 WB ;Rat. 38726273
- [IF=4.6] Demircan Volkan. et al. Evaluation of therapeutic use of a combination of pentoxifylline and vitamin E in radiation-induced renal fibrosis. SCI REP-UK. 2024 Mar;14(1):1-14 IHC; Rat. 38521858