

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Mafa Rabbit pAb

Catalog Number: bs-22235R

Target Protein: Mafa
Concentration: 1mg/ml

Form: Liquid
Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Mouse, Rat (predicted:Human, Pig, Cow, Chicken)

Predicted MW: 37 kDa
Entrez Gene: 389692
Swiss Prot: Q8NHW3

Source: KLH conjugated synthetic peptide derived from human Mafa: 71-170/352.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: Insulin gene expression is regulated by several islet-enriched transcription factors. However,

MAFA is the only beta cell-specific activator. MAFA selectively induces endogenous insulin transcription in non-beta cells. MAFA was also first detected in the insulin-producing cells formed during the second and predominant phase of beta cell differentiation, and absent in the few insulin-positive cells found in Nkx6.1(-/-) pancreata, which lack the majority of second-phase beta cells. These results demonstrate that MAFA is a potent insulin activator

that is likely to function downstream of Nkx6.1 during islet insulin-producing cell

development.

VALIDATION IMAGES



Sample: Cerebrum (Rat) Lysate at 40 ug Lung (Mouse) Lysate at 40 ug Spleen (Mouse) Lysate at 40 ug Primary: Anti- Mafa (bs-22235R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 37 kD Observed band size: 50 kD

PRODUCT SPECIFIC PUBLICATIONS

[IF=5] Peibin Lin. et al. Naringenin protects pancreatic β cells in diabetic rat through activation of estrogen receptor β . EUR J PHARMACOL. 2023 Dec;960:176115 WB; Rat . 37866740