bs-13153R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

FDPS Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 2224 SWISS: P14324

Target: FDPS

Immunogen: KLH conjugated synthetic peptide derived from human FDPS:

201-300/419.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: FDPS is a 419 amino acid enzyme belonging to the FPP/GGPP

synthetase family. Localized to cytoplasm and peroxisome, FDPS expression is regulated by phorbol esters and polyunsaturated fatty acids. FDPS assists in cholesterol biosynthesis, posttranslational protein modifications and synthesis of steroid hormones in the isoprenoid pathway.FDPS catalyzes the formation of farnesyl diphosphate (FPP), a precursor for several classes of essential metabolites including sterols, dolichols, carotenoids, and ubiquinones. FDPS is inactivated by interferon-induced RSAD2, which may result in the disruption of lipid rafts at the plasma membrane. Existing as a homodimer, FDPS may have anti-viral effects when inactivated by RSAD2. Reduced activity of FDPS in liver may partly be the cause of Zellweger syndrome and neonatal adrenoleukodystrophy, both of which are known to be

peroxisomal deficiency diseases.

Applications: WB (1:500-2000)


IHC-P (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500) **ELISA** (1:5000-10000)

Reactivity: Human, Mouse, Rat


Predicted 48 kDa MW.:

Subcellular Location: Cytoplasm


VALIDATION IMAGES



Sample: Lane 1: Mouse Liver tissue lysates Lane 2: Mouse Kidney tissue lysates Lane 3: Rat Liver tissue lysates Lane 4: Rat Kidney tissue lysates Lane 5: Human HepG2 cell lysates Lane 6: Human HeLa cell lysates Primary: Anti-FDPS (bs-13153R) at 1/500 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 48 kDa Observed band size: 40 kDa



Paraformaldehyde-fixed, paraffin embedded (human liver carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (FDPS) Polyclonal Antibody, Unconjugated (bs-13153R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (FDPS) Polyclonal Antibody, Unconjugated (bs-13153R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.