

**bsm-51134M****[ Primary Antibody ]****GABARAP Mouse mAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**

<b>Host:</b> Mouse	<b>Isotype:</b> IgG1	<b>Applications:</b> <b>WB</b> (1:500-2000) <b>ELISA</b> (1:5000-10000)
<b>Clonality:</b> Monoclonal	<b>CloneNo.:</b> 10B2	
<b>GeneID:</b> 11337	<b>SWISS:</b> O95166	
<b>Target:</b> GABARAP		
<b>Purification:</b> affinity purified by Protein G		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		<b>Reactivity:</b> Human, Mouse (predicted: Rat, Rabbit, Cow)
<b>Background:</b> Gamma-aminobutyric acid A receptors [GABA(A) receptors] are ligand-gated chloride channels that mediate inhibitory neurotransmission. This gene encodes GABA(A) receptor-associated protein, which is highly positively charged in its N-terminus and shares sequence similarity with light chain-3 of microtubule-associated proteins 1A and 1B. This protein clusters neurotransmitter receptors by mediating interaction with the cytoskeleton. [provided by RefSeq, Jul 2008]		<b>Predicted MW.:</b> 14 kDa
		<b>Subcellular Location:</b> Cell membrane ,Cytoplasm

**— VALIDATION IMAGES —**

Anti-GABARAP Antibody at 1:1000 dilution + mouse brain lysate Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 13918 Da  
Blocking/Dilution buffer: 5% NFDm/TBST.



Anti- at 1:1000 dilution + SH-SY5Y whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 14 kDa Blocking/Dilution buffer: 5% NFDm/TBST.