

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## **Recombinant human ACAN protein, N-His**

Concentration: >0.5 mg/ml AA Seq: 2321-2530/2530 Predicted MW: 27.28 Detected MW: 30 kDa Tags: N-His Activity: Not tested
Predicted MW: 27.28 Detected MW: 30 kDa Tags: N-His
Detected MW: 30 kDa Tags: N-His
Tags: N-His
Activity: Not tested
Activity. Horizotta
Endotoxin: Not analyzed
Purity: >95% as determined by SDS-PAGE
Purification: AC
Form: Lyophilized or Liquid
Storage: PBS (pH7.4).
Stored at -70°C or -20°C. Avoid repeated freeze/thaw cycles.
Background: Aggrecan is a member of a family of large, aggregating proteoglycans (also including
versican, brevican and neurocan) which is found in articular cartilage. Aggrecan is
composed of three major domains: G1, G2, and G3. Between the G1 and G2 domains there i
an interglobulin region (IGD). The IGD region is the major site of cleavage by specific
proteases like metalloproteinases (MMPs) and aggrecanase. Aggrecan cleavage has been
associated with a number of degenerative diseases including rheumatoid arthritis and
osteoarthritis. There is evidence that this family of proteoglycans modulates cell adhesion,
migration, and axonal outgrowth in the CNS.

## VALIDATION IMAGES



The purity of the protein is greater than 90% as determined by reducing SDS-PAGE.

The purity of the protein is greater than 90% as determined by reducing SDS-PAGE.