## bsm-52028M

# [ Primary Antibody ]

# Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# **Cdk4 Mouse mAb**

- DATASHEET -

Host: Mouse Isotype: IgG
Clonality: Monoclonal CloneNo.: 2F6
GeneID: 1019 SWISS: P11802

Target: Cdk4

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** The protein encoded by this gene is a member of the Ser/Thr

protein kinase family. This protein is highly similar to the gene products of S. cerevisiae cdc28 and S. pombe cdc2. It is a catalytic subunit of the protein kinase complex that is important for cell cycle G1 phase progression. The activity of this kinase is restricted to the G1-S phase, which is controlled by the regulatory subunits D-type cyclins and CDK inhibitor p16(INK4a). This kinase was shown to be responsible for the phosphorylation of retinoblastoma gene product (Rb). Mutations in this gene as well as in its related proteins including D-type cyclins, p16(INK4a) and Rb were all found to be associated with tumorigenesis of a variety of cancers. Multiple polyadenylation sites of this gene have been reported. [provided by RefSeq, Jul 2008]

Applications: WB (1:500-2000)


IHC-P (1:200-1000) IHC-F (1:200-1000) IF (1:200-1000) ICC/IF (1:50-200)

Reactivity: Human, Mouse, Rat


Predicted MW.: 34 kDa

**Subcellular** Cell membrane ,Cytoplasm **Location:** ,Nucleus


### VALIDATION IMAGES



Sample: Lane 1: Embryo (Mouse) Lysate at 40 ug Lane 2: Placenta (Mouse) Lysate at 40 ug Lane 3: Heart (Mouse) Lysate at 40 ug Lane 3: Heart (Mouse) Lysate at 40 ug Lane 4: NIH/3T3 (Mouse) Cell Lysate at 30 ug Lane 5: Heart (Rat) Lysate at 40 ug Lane 6: Cerebrum (Rat) Lysate at 40 ug Lane 7: Jurkat (Human) Cell Lysate at 30 ug Lane 8: Hela (Human) Cell Lysate at 30 ug Lane 9: MCF-7 (Human) Cell Lysate at 30 ug Lane 10: A431 (Human) Cell Lysate at 30 ug Primary: Anti-Cdk4 (bsm-52028M) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution Predicted band size: 34 kD Observed band size: 32 kD



Paraformaldehyde-fixed, paraffin embedded (human rectal carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Cdk4) Monoclonal Antibody, Unconjugated (bsm-52028M) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human myocardium); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Cdk4) Monoclonal Antibody, Unconjugated (bsm-52028M) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.

### - SELECTED CITATIONS -

- [IF=11.205] Cong Lan. et al. Inhibition of DYRK1A, via histone modification, promotes cardiomyocyte cell cycle activation and cardiac repair after myocardial infarction. EBIOMEDICINE. 2022 Aug;82:104139 WB;Rat. 35810562
- [IF=10.75] Tian Xiangdong. et al. Autophagy Inhibition Contributes to Apoptosis of PLK4 Downregulation-induced Dormant Cells in Colorectal Cancer. INT J BIOL SCI. 2023 May;19(9):2817-2834 WB; Human. 37324947

- [IF=8.469] Que, Tianshi. et al. HMGA1 stimulates MYH9-dependent ubiquitination of GSK-3ß via PI3K/Akt/c-Jun signaling to promote malignant progression and chemoresistance in gliomas. Cell Death Dis. 2021 Dec;12(12):1-12 WB; Human. 34887392
- [IF=8.2] Chun Huang. et al. Whole-transcriptome analysis of longissimus dorsi muscle in cattle-yaks reveals the regulatory functions of ADAMTS6 gene in myoblasts. INT J BIOL MACROMOL. 2024 Mar;262:129985 WB; Mouse. 38342263
- [IF=4.5] Dong-Dong Wang. et al. Identification of diterpenoids from Salvia castanea Diels f. tomentosa Stib and their antitumor activities. BIOORG CHEM. 2024 Aug;:107701 WB; Human. 39154520