

---

## NTN3 Rabbit pAb

Catalog Number: bs-11059R

Target Protein: NTN3

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: **IHC-P** (1:100-500), **IHC-F** (1:100-500), **IF** (1:100-500)

Reactivity: Human, Rat (predicted:&nbsp;Mouse, Pig, Sheep, Cow, Dog)

Predicted MW: 59 kDa

Subcellular: Secreted ,Extracellular matrix

Locations:

Entrez Gene: 4917

Swiss Prot: O00634

Source: KLH conjugated synthetic peptide derived from human NTN3/Netrin-3: 401-500/580.

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


**Background:** Netrin proteins are a family of laminin-related secreted proteins that provide guidance signals for axonal growth and cell migration during development. Netrin signaling is dependent on the concentration of calcium outside the cell and the level of PKA activity. In axonal cells, a reduction in PKA activity converts the responsiveness of the axons to the netrin proteins as the cells are repelled, rather than attracted, by the netrin gradient. Neogenin serves as the primary guidance receptor for netrin-3. Netrin-2 and the corresponding mouse homolog netrin-3 are expressed primarily in the lower two-thirds of the spinal cord, and, like netrin-1, they can either attract or repel commissural axons at a distance. Netrin-3 proteins are associated with the axon fibers projecting from motor neurons and from neurons within sympathetic and sensory ganglia, suggesting that netrin-3 may be involved in pathfinding and fasciculation of axon projection. Neogenin serves as the primary guidance receptor for netrin-3. During peripheral nerve development, high netrin-3 expression has been detected in mesenchymal cells, sensory ganglia and muscles. In humans, the gene encoding for the netrin-3 protein is localized to chromosome 16p13.3.

## VALIDATION IMAGES

---



Paraformaldehyde-fixed, paraffin embedded (Rat spinal cord); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NTN3) Polyclonal Antibody, Unconjugated (bs-11059R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human skin carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NTN3) Polyclonal Antibody, Unconjugated (bs-11059R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.