

bsm-60761R**[Primary Antibody]****CD206 Recombinant Rabbit mAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:200-1000) IHC-F (1:200-1000) IF (1:200-1000) ICC/IF (1:50-200) Reactivity: Human Predicted MW.: 160 kDa Subcellular Location: Cell membrane
Clonality: Recombinant	CloneNo.: 12A3	
GeneID: 4360	SWISS: P22897	
Target: CD206		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: PBS, Glycerol, BSA. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The recognition of complex carbohydrate structures on glycoproteins is an important part of several biological processes, including cell-cell recognition, serum glycoprotein turnover, and neutralization of pathogens. The protein encoded by this gene is a type I membrane receptor that mediates the endocytosis of glycoproteins by macrophages. The protein has been shown to bind high-mannose structures on the surface of potentially pathogenic viruses, bacteria, and fungi so that they can be neutralized by phagocytic engulfment. This gene is in close proximity to MRC1L1. The gene loci including this gene, MRC1L1, as well as LOC340843 and LOC340893, consist of two nearly identical, tandemly linked genomic regions, which are thought to be a part of a duplicated region. [provided by RefSeq].		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded Human Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD206 Monoclonal Antibody, Unconjugated(bsm-60761R) at 1:1000 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023)and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD206 Monoclonal Antibody, Unconjugated(bsm-60761R) at 1:1000 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023)and DAB (C-0010) staining.



4% Paraformaldehyde-fixed Molt-4 (H) cell; Triton X-100 at r.t. for 20 min; Antibody incubation with (MRC1) monoclonal Antibody, unconjugated (bsm-60761R) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-60295G-FITC) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.

— SELECTED CITATIONS —

- **[IF=13.3]** Zhihong Su. et al. Novel asymmetrical double-layer structural adhesive hydrogels with synergetic neuroprotection and angiogenesis effect for diabetic wound healing. CHEM ENG J. 2024 Dec;;159081 IF ;Mouse. 10.1016/j.cej.2024.159081
- **[IF=13.3]** Lubin Zhou. et al. Electrospun Self-Pumping dressing with gastrodin for immunomodulation and rapid healing of diabetic wounds. CHEM ENG J. 2024 Sep;495:153424 IF ;Rat. 10.1016/j.cej.2024.153424
- **[IF=11]** Ou Da-Liang. et al. Nano-modified viruses prime the tumor microenvironment and promote the photodynamic virotherapy in liver cancer. J BIOMED SCI. 2024 Dec;31(1):1-12 IHC ;Mouse. 38163894

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

- **[IF=10]** Lubin Zhou. et al. A Novel Self-Pumping Janus Dressing for Promoting Wound Immunomodulation and Diabetic Wound Healing. ADV HEALTHC MATER. 2023 Nov;;2303460 IF ;Rat. 37957786
- **[IF=8.7]** Xiaolan Ou. et al. Portable direct spraying porous nanofibrous membranes stent-loaded polymyxin B for treating diabetic wounds with difficult-to-heal gram-negative bacterial infections. MATER TODAY BIO. 2024 Dec;29:101365 IF ;Rat. 39687800