
bs-22547R	[ Primary Antibody ]	Bioss	
hydrogen exchan	iger 3 Rabbit pAb	ANTIBODIES www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800	
Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)	
Clonality: Polyclonal	,per 1g0	, , , , , , , , , , , , , , , , , , ,	
GenelD: 6550	SWISS: P48764	<b>Reactivity:</b> Mouse, Rat (predicted: Human, Rabbit,	
Target: hydrogen excha	nger 3	Pig, Sheep, Cow)	
Immunogen: KLH conjugated synthetic peptide derived from human hydrogen exchanger 3 : 581-680/834. < Cytoplasmic >		Predicted <sub>93 kDa</sub>	
Purification: affinity purified by Protein A		Subsollular	
Concentration: 1mg/ml		Subcellular Location: Cell membrane	
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.			
<b>Background:</b> NHE-3 are integral membrane proteins that are expressed in most mammalian tissues, where they regulate intracellular pH and cell volume. NHEs mediate the transport of hydrogen (H+) ions out of cells in exchange for extracellular sodium (Na+) ions. While NHE-1 is ubiquitously expressed, the NHE isoforms 2-8 have distinct tissue- and cell type-dependent expression and inhibitory characteristics. NHE-3 localizes to the apical membrane of renal proximal tubules where it is responsible for most of the sodium transport and fluid reabsorption. NHE-3 translocates to internal pools where it mediates natriuresis when blood pressure increases abruptly. NHE-3 is also expressed in the stomach and functions to protect the mucosa by secreting protons that diffuse into the mucous cells.			

## - VALIDATION IMAGES -



Sample: Kidney (Mouse) Lysate at 40 ug Kidney (Rat) Lysate at 40 ug Primary: Anti-hydrogen exchanger 3 (bs-22547R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 80 kD Observed band size: 80 kD