

bs-24579R**[Primary Antibody]****NOBOX Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:400-800) IHC-F (1:400-800) IF (1:100-500)
Clonality: Polyclonal		Reactivity: Rat (predicted: Mouse)
Target: NOBOX		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: 57 kDa
Background: Early ovarian folliculogenesis is characterized by the breakdown of germ cell clusters and formation of primordial follicles. The cessation of ovarian function under the age of 40 years results in premature ovarian failure (POF) and is accompanied by amenorrhea, hypoeestrogenism and elevated serum gonadotropin concentrations. 1% of all women are affected by POF, and mutations in a few genes, including inhibin β fsh receptor and the LH/choriogonadotropin receptor have been linked to POF. In addition, several germ cell specific genes and downstream transcription factors are thought to play an important role in human oogenesis. NOBOX (newborn ovary homeobox gene), an oocyte-specific homeobox gene, is a critical protein involved in early folliculogenesis. Missense mutations have been shown to cause nonsyndromic ovarian failure by disrupting the NOBOX protein's ability to bind to NOBOX DNA-binding element (NBE), and thereby inhibiting its regulation of Pou5f1 and GDF-9. NOBOX expression in the ovary is oocyte specific, but it shows expression in adult testis and pancreas as well.		Subcellular Location: Nucleus

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (rat uterus); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NOBOX) Polyclonal Antibody, Unconjugated (bs-24579R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=6.055]** Juan Feng. et al. Melatonin prevents cyclophosphamide-induced primordial follicle loss by inhibiting ovarian granulosa cell apoptosis and maintaining AMH expression. FRONT ENDOCRINOL. 2022; 13: 895095 IHC ;Mouse. 35992124

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.