

bs-11773R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Transglutaminase 6 Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Rat (predicted: Human, Mouse, Sheep, Cow, Dog, Horse) Predicted MW.: 79 kDa Subcellular Location: Cytoplasm
Clonality: Polyclonal		
GeneID: 343641	SWISS: Q95932	
Target: Transglutaminase 6		
Immunogen: KLH conjugated synthetic peptide derived from human TGase6/Transglutaminase 6: 551-640/706.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Terminally differentiating mammalian epidermal cells acquire an insoluble, 10 to 20 nm thick protein deposit on the intracellular surface of the plasma membrane, known as the cross-linked cell envelope (CE). The CE is a component of the epidermis that is generated through the formation of disulfide bonds and gamma-glutamyl-lysine isodipeptide bonds, which are formed by the action of transglutaminases (TGases). TGases are Ca ²⁺ -dependent enzymes, which catalyze the formation of isopeptide bonds by transferring an amine to a glutamyl residue, thereby cross-linking glutamine residues and lysine residues in substrate proteins. TGases influence numerous biological processes, including blood coagulation, epidermal differentiation, seminal fluid coagulation, fertilization, cell differentiation and apoptosis. TGase6 (transglutaminase 6), also known as TGM6, TGY or TGM3L, is a 706 amino acid protein that catalyzes the cross-linking of proteins and the conjugation of proteins to polyamines. As a result of alternative splicing, two TGase6 isoforms exist.		

— VALIDATION IMAGES —

Tissue/cell: Rat kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-TGM6 Polyclonal Antibody, Unconjugated(bs-11773R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining