

**bs-9582R****[ Primary Antibody ]****phospho-NHE3 (Ser552) Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<p><b>Host:</b> Rabbit</p> <p><b>Clonality:</b> Polyclonal</p> <p><b>GeneID:</b> 6550</p> <p><b>Target:</b> phospho-NHE3 (Ser552)</p> <p><b>Immunogen:</b> KLH conjugated synthesised phosphopeptide derived from rat NHE-3 around the phosphorylation site of Ser552 (human Ser555): RG(p-S)LA.</p> <p><b>Purification:</b> affinity purified by Protein A</p> <p><b>Concentration:</b> 1mg/ml</p> <p><b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p><b>Background:</b> NHE-3 are integral membrane proteins that are expressed in most mammalian tissues, where they regulate intracellular pH and cell volume. NHEs mediate the transport of hydrogen (H<sup>+</sup>) ions out of cells in exchange for extracellular sodium (Na<sup>+</sup>) ions. While NHE-1 is ubiquitously expressed, the NHE isoforms 2-8 have distinct tissue- and cell type-dependent expression and inhibitory characteristics. NHE-3 localizes to the apical membrane of renal proximal tubules where it is responsible for most of the sodium transport and fluid reabsorption. NHE-3 translocates to internal pools where it mediates natriuresis when blood pressure increases abruptly. NHE-3 is also expressed in the stomach and functions to protect the mucosa by secreting protons that diffuse into the mucous cells.</p>	<p><b>Isotype:</b> IgG</p> <p><b>SWISS:</b> P48764</p> <p><b>Applications:</b> <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>ICC/IF</b> (1:100-500) <b>ELISA</b> (1:5000-10000)</p> <p><b>Reactivity:</b> (predicted: Human, Mouse, Rat, Rabbit, Pig, Cow, Chicken, Dog, Horse)</p> <p><b>Predicted MW.:</b> 93 kDa</p> <p><b>Subcellular Location:</b> Cell membrane</p>
--	--