

**bs-2977R****[ Primary Antibody ]****BioSS**  
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**TAGL2 Rabbit pAb****— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> IHC-P (1:100-500)
<b>Clonality:</b> Polyclonal		<b>IHC-F</b> (1:100-500)
<b>GeneID:</b> 8407	<b>SWISS:</b> P37802	<b>IF</b> (1:100-500)
<b>Target:</b> TAGL2		<b>Flow-Cyt</b> (2ug/Test)
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human TAGL2: 51-160/199.		<b>Reactivity:</b> Human, Mouse (predicted: Rat, Pig, Cow, Dog)
<b>Purification:</b> affinity purified by Protein A		<b>Predicted MW.:</b> 22 kDa
<b>Concentration:</b> 1mg/ml		<b>Subcellular Location:</b> Secreted ,Extracellular matrix ,Cytoplasm
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> The protein encoded by this gene is similar to the protein transgelin, which is one of the earliest markers of differentiated smooth muscle. The specific function of this protein has not yet been determined, although it is thought to be a tumor suppressor. Multiple transcript variants encoding different isoforms have been found for this gene.		

**— VALIDATION IMAGES —**

Tissue/cell: muscle of mouse embryo; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-TAGL2 Polyclonal Antibody, Unconjugated(bs-2977R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Blank control: A549(blue), the cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice.. Isotype Control Antibody: Rabbit IgG(orange) ; Secondary Antibody: Goat anti-rabbit IgG-FITC(white blue), Dilution: 1:100 in 1 X PBS containing 0.5% BSA ; Primary Antibody Dilution: 5µg in 100 µL 1X PBS containing 0.5% BSA(green).