

bs-20101R**[Primary Antibody]**

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Angiotensin III Rabbit pAb


— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1ug/Test)
Clonality: Polyclonal		
GeneID: 183	SWISS: P01019	
Target: Angiotensin III		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Reactivity: Human, Mouse, Rat (predicted: Horse)
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: bs-0587P is a Eight branched multiple antigenic peptide of Angiotensin II.		
		Predicted MW.: 1/50 kDa Subcellular Location: Secreted


— VALIDATION IMAGES —




Sample: Lane 1: Liver (Mouse) Lysate at 40 ug
 Lane 2: Plasma (Mouse) at 20 ug Lane 3: Kidney (Mouse) Lysate at 40 ug Lane 4: Adrenal gland (Mouse) Lysate at 40 ug Lane 5: Adrenal gland (Rat) Lysate at 40 ug Lane 6: HepG2 (Human) Cell Lysate at 30 ug
 Primary: Anti-Angiotensin III (bs-20101R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 52/60 kD Observed band size: 52/60 kD



Paraformaldehyde-fixed, paraffin embedded (Rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Angiotensin III) Polyclonal Antibody, Unconjugated (bs-20101R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Angiotensin III) Polyclonal Antibody, Unconjugated (bs-20101R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control: HepG2. Primary Antibody (green line): Rabbit Anti-Angiotensin III antibody (bs-20101R) Dilution: 1μg / 10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: 1μg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 0.1% PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=3.367]** Hua Chen. et al. Chronic Intermittent Hypobaric Hypoxia Decreases High Blood Pressure by Stabilizing the Vascular Renin-Angiotensin System in Spontaneously Hypertensive Rats. Front Physiol. 2021; 12: 639454 IHC ;Rat. 33841179
- **[IF=3.2]** Delong Duo. et al. Long-term exposure to high-altitude hypoxic environments reduces blood pressure by inhibiting the renin-angiotensin system in rats. FRONT PHYSIOL. 2025 Apr;16:1565147 WB ;Rat. 40303590