### bs-10136R

## [ Primary Antibody ]

# Caspase-14 p10 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

		400-901-9800
Host: Rabbit	Isotype:  gG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		<b>IHC-F</b> (1:100-500)
GenelD: 23581	SWISS: P31944	
Target: Caspase-14 p10		<b>Reactivity:</b> Human, Mouse, Rat (predicted: GuineaPig)
Immunogen: KLH conjugated sy subunit p10: 155-2	nthetic peptide derived from human Caspase-14 42/242.	
Purification: affinity purified by Protein A		Predicted
Concentration: 1mg/ml		MVV.:
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: <sup>Cytoplasm</sup>
<b>Background:</b> Believed to be a no epidermal differen differentiation and the epidermis by p	on-apoptotic caspase which is involved in tiation. Seems to play a role in keratinocyte I cornification. Probably regulates maturation of roteolytically processing filaggrin.	

#### VALIDATION IMAGES



Tissue/cell: rat colon tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-Caspase-14 Polyclonal Antibody, Unconjugated(bs-10136R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-caspase-14 Polyclonal Antibody, Unconjugated(bs-10136R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

### - SELECTED CITATIONS -

• [IF=6.8] Zhang Hongrong. et al. Multi-omics analysis deciphers intercellular communication regulating oxidative stress to promote oral squamous cell carcinoma progression. NPJ PRECIS ONCOL. 2024 Nov;8(1):1-18 IF ;Human. 39572698