

bs-8611R**[Primary Antibody]****USP22 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 23326**SWISS:** Q9UPT9**Target:** USP22**Immunogen:** KLH conjugated synthetic peptide derived from human USP22: 121-220/525.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The ubiquitin (Ub) pathway involves three sequential enzymatic steps that facilitate the conjugation of Ub and Ub-like molecules to specific protein substrates. Through the use of a wide range of enzymes that can add or remove ubiquitin, the Ub pathway controls many intracellular processes such as signal transduction, transcriptional activation and cell cycle progression. USP22 (ubiquitin specific peptidase 22), also known as USP3L, is a 525 amino acid protein that contains one UBP-type zinc finger and functions to catalyze the conversion of a ubiquitin C-terminal thioester to free ubiquitin and thiol, a reaction that may influence several cellular processes. Via its catalytic activity, USP22 is thought to play an important role in cell cycle progression and may also serve as a cancer stem cell marker.**Applications:** **WB** (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Reactivity:** Human, Rat
(predicted: Mouse, Rabbit, Sheep, Cow, Chicken, Dog, Horse, Xenopus laevis)**Predicted MW.:** 60 kDa**Subcellular Location:** Nucleus**— VALIDATION IMAGES —**

Sample: Lane 1: Human HeLa cell lysates Lane 2: Human A549 cell lysates Lane 3: Human Jurkat cell lysates Lane 4: Human 293T cell lysates
 Primary: Anti-USP22 (bs-8611R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 60 kDa Observed band size: 60 kDa



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-USP22 Polyclonal Antibody, Unconjugated(bs-8611R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining