

bsm-63132R

[Primary Antibody]

Mono-Methyl-Histone H4 (K17) Recombinant Rabbit mAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:1000-1:2000) ICC/IF (1:50-1:200) Reactivity: Human, Mouse Predicted MW.: 11 kDa Subcellular Location: Nucleus
Clonality: Recombinant	CloneNo.: 4G1	
GeneID: 121504	SWISS: P62805	
Target: Mono-Methyl-Histone H4 (K17)		
Immunogen: A synthesized peptide derived from human Histone H4 around the methylation site of K17: GA-K(Me)-RH(全长103aa).		
Purification: affinity purified by Protein A		
Storage: 10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid repeated freeze/thaw cycles.		
Background: Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.		