## bsm-63132R

- DATASHEET ------

## [ Primary Antibody ]

## Mono-Methyl-Histone H4 (K17) Recombinant Rabbit mAb



sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Host: Rabbit	<b>Isotype:</b> IgG	Applications: WB (1:1000-2000)
Clonality: Recombinant	CloneNo.: 4G1	ICC/IF (1:50-200)
GeneID: 121504	SWISS: P62805	Reactivity: Human, Mouse
Target: Mono-Methyl-Histone H	14 (K17)	
<b>Immunogen:</b> A synthesized peptide derived from human Histone H4 around the methylation site of K17: GA-K(Me)-RH(全长103aa).		Predicted 11 kDa MW.:
Purification: affinity purified by Protein A		
<b>Storage:</b> 10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid repeated freeze/thaw cycles.		Subcellular Location: Nucleus
<b>Background:</b> Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.		