bsm-63150R

[Primary Antibody]

Bioss ANTIBODIES

phospho-BTK (Y551) Recombinant Rabbit mAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG
Clonality: Recombinant CloneNo.: 5C12
GeneID: 695 SWISS: Q06187

Target: BTK (Y551)

Immunogen: A synthesized peptide derived from human BTK around the

phosphorylation site of Y551: DE-pY-TSS(全长659aa).

Purification: affinity purified by Protein A

Storage: 10mM phosphate buffered saline(pH 7.4) with 150mM sodium

chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid

repeated freeze/thaw cycles.

Background: Non-receptor tyrosine kinase indispensable for B lymphocyte

development, differentiation and signaling. Binding of antigen to the B-cell antigen receptor (BCR) triggers signaling that ultimately leads to B-cell activation. After BCR engagement and activation at the plasma membrane, phosphorylates PLCG2 at several sites, igniting the downstream signaling pathway through calcium mobilization, followed by activation of the protein kinase C (PKC) family members.

Applications: WB (1:1000-2000)

ICC/IF (1:50-200)

Reactivity: Human

Predicted MW.: 76 kDa

Subcellular Cell membrane, Cytoplasm

Location: , Nucleus