## bsm-63153R

## [ Primary Antibody ]

## Histone H2B (Acetyl K20) Recombinant Rabbit mAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: WB (1:1000-5000)
Clonality: Recombinant	CloneNo.: 8D1	IHC-P (1:100-200) IHC-F (1:100-200)
GenelD: 3018	SWISS: P33778	<b>IF</b> (1:50-200)
<b>Target:</b> Histone H2B (Acetyl K20) <b>Immunogen:</b> A synthesized peptide derived from human Histone H2B around the acetylation site of K20: VT-(acetyl)K-AQ.		Flow-Cyt (1:50-100) ICC/IF (1:50-200) IP (1:20-50) ChIP (1:20-50)
<ul> <li>Purification: affinity purified by Protein A</li> <li>Storage: 10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid repeated freeze/thaw cycles.</li> </ul>		Reactivity: Human, Mouse, Rat Predicted MW.: <sup>14</sup> kDa Subcellular Location: <sup>Nucleus</sup>
<b>Background:</b> Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.		