

bsm-63169R

[Primary Antibody]

HDAC10 Recombinant Rabbit mAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit

Isotype: IgG

Clonality: Recombinant

CloneNo.: 3F3

GeneID: 83933

SWISS: Q969S8

Target: HDAC10

Immunogen: A synthesized peptide derived from human HDAC10: 345-383.

Purification: affinity purified by Protein A

Storage: 10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid repeated freeze/thaw cycles.

Background: Polyamine deacetylase (PDAC), which acts preferentially on N8-acetylspermidine, and also on acetylcadaverine and acetylputrescin. Exhibits attenuated catalytic activity toward N1,N8-diacetylspermidine and very low activity, if any, toward N1-acetylspermidine.

Applications: WB (1:1000-1:2000)

ICC/IF (1:50-1:200)

IP (1:20-1:50)

Reactivity: Human, Mouse

Predicted

MW.: 71 kDa

Subcellular

Location: Cytoplasm ,Nucleus