bsm-63244R

[Primary Antibody]

Acetyl-Histone H3(K10) Recombinant Rabbit mAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG
Clonality: Recombinant CloneNo.: 3A7
GeneID: 8350 SWISS: P68431

Target: Acetyl-Histone H3(K10)

Immunogen: A synthesized peptide derived from human Histone H3.1 around the

acetylation site of K10: TAR-K(Ac)-STG.

Purification: affinity purified by Protein A

Storage: 10mM phosphate buffered saline(pH 7.4) with 150mM sodium

chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid

repeated freeze/thaw cycles.

Background: Core component of nucleosome. Nucleosomes wrap and compact

DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also

called histone code, and nucleosome remodeling.

Applications: WB (1:1000-1:2000)

IHC-P (1:100-1:200) IHC-F (1:100-1:200) IF (1:50-1:200) Flow-Cyt (1:20-1:100) ICC/IF (1:50-1:200)

Reactivity: Human, Mouse, Rat

IP (1:20-1:50)

Predicted MW.: 15 kDa

Subcellular Nucleus