

bsm-63270R

[Primary Antibody]

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

HSF1 Recombinant Rabbit mAb

DATASHEET

Host: Rabbit

Clonality: Recombinant

GeneID: 3297

Target: HSF1

Immunogen: A synthesized peptide derived from human HSF1: 485-529.

Purification: affinity purified by Protein A

Storage: 10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid repeated freeze/thaw cycles.

Background: Functions as a stress-inducible and DNA-binding transcription factor that plays a central role in the transcriptional activation of the heat shock response (HSR), leading to the expression of a large class of molecular chaperones, heat shock proteins (HSPs), that protect cells from cellular insult damage.

Isotype: IgG

CloneNo.: 4F3

SWISS: Q00613

Applications: WB (1:500-1:2000)

IHC-P (1:100-1:500)

IHC-F (1:100-1:500)

IF (1:100-1:500)

Flow-Cyt (1µg/Test)

ICC/IF (1:50-1:200)


IP (1:20-1:50)

Reactivity: Human


Predicted
MW.: 57 kDa

Subcellular
Location: Cytoplasm ,Nucleus


VALIDATION IMAGES




25 ug total protein per lane of various lysates (see on figure) probed with HSF1 monoclonal antibody, unconjugated (bsm-63270R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Paraformaldehyde-fixed, paraffin embedded Human Breast Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with HSF1 Monoclonal Antibody, Unconjugated(bsm-63270R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human ovary; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with HSF1 Monoclonal Antibody, Unconjugated(bsm-63270R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



4% Paraformaldehyde-fixed HeLa (H) cell; Triton X-100 at r.t. for 20 min; Antibody incubation with (HSF1) monoclonal Antibody, unconjugated (bsm-63270R) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-40295G-FITC) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.



The HeLa (H) cells were fixed with 4% PFA (10 min at r.t.) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, the cells then were incubated in 5%BSA to block non-specific protein-protein interactions (30 min at r.t.). Primary Antibody (green): Rabbit Anti-HSF1 antibody (bsm-63270R): 1:50-100/10⁶ cells; Secondary Antibody (white blue): Goat anti-Rabbit IgG-BF488 (bs-60295G-BF488): 1 µg/test. Isotype Control (orange): Rabbit IgG (bs-0295P). Blank control (black): PBS. Acquisition of 20,000 events was performed.