

---

## GGA2 Rabbit pAb

Catalog Number: bs-13344R

Target Protein: GGA2

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Rat (predicted:Human, Mouse, Rabbit, Pig, Sheep, Cow, Dog)

Predicted MW: 67 kDa

Subcellular: Cell membrane ,Cytoplasm

Locations:

Entrez Gene: 23062

Swiss Prot: Q9UJY4

Source: KLH conjugated synthetic peptide derived from human GGA2: 161-260/613.

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


**Background:** A family of proteins, the GGAs (Golgi-localized, g-adaptin ear-containing, ARF-binding proteins) sequences that showed significant homology to the carboxy-terminal 'ear' domain of g-adaptin. Members of the GGA family (GGA1, GGA2 (also known as VEAR or VHS domain and ear domain of g-adaptin) and GGA3) are ubiquitous coat proteins that facilitate the trafficking of proteins between the trans-Golgi network and the lysosome. However, unlike g-adaptin, the GGAs are not associated with clathrin-coated vesicles or with any of the components of the AP-1 complex. GGA1 and GGA2 are also not associated with each other, although they colocalize on perinuclear membranes. GGA2 shares 45% amino acid sequence identity with GGA1 and 35% with GGA3. In addition to being involved in heterotypic vesicle/suborganelle interactions associated with the Golgi complex, GGA2 may have a tissue-specific function and is highly expressed in kidney, muscle and heart. Furthermore, the VHS domain of GGA2 binds to the acidic cluster-di-leucine motif in the cytoplasmic tail of the cation-independent mannose 6-phosphate receptor (CI-MPR) and this is important for lysosomal enzyme targeting.

## VALIDATION IMAGES


---



Tissue/cell: Rat brain; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-GGA2, Unconjugated(bs-13344R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-GGA2 Polyclonal Antibody, Unconjugated(bs-13344R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-GGA2 Polyclonal Antibody, Unconjugated(bs-13344R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining