bs-6936R

[Primary Antibody]

MYF5 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:50-200)

Reactivity: Mouse, Rat (predicted: Human, Rabbit, Pig, Sheep, Cow, Chicken, Horse)

Predicted MW.: ^{28 kDa}

Subcellular Location: Nucleus

Host: Rabbit

- DATASHEET -

Clonality: Polyclonal GeneID: 4617

SWISS: P13349

Isotype: IgG

Target: MYF5

Immunogen: KLH conjugated synthetic peptide derived from human MYF5: 61-160/255.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Differentiation of myogenic cells is regulated by multiple positively and negatively acting factors. One well characterized family of helix-loop-helix (HLH) proteins known to play an important role in the regulation of muscle cell development include Myo D, myogenin, Myf-5 and Myf-6 (also designated MRF-4 or herculin). Of interest, most muscle cells express either Myo D or Myf-5 in the committed state, but when induced to differentiate, all turn on expression of myogenin. Myo D transcription factors form heterodimers with products of a more widely expressed family of bHLH genes, the E family, which consists of at least three distinct genes: E2A, IF2 and HEB. Myo D-E heterodimers bind avidly to consensus (CANNTG) E box target sites that are functionally important elements in the upstream regulatory sequences of many muscle-specific terminal differentiation genes.

– VALIDATION IMAGES



Sample: Lane 1: Muscle (Mouse) Lysate at 40 ug Lane 2: Tongue (Mouse) Lysate at 40 ug Lane 3: Muscle (Rat) Lysate at 40 ug Lane 4: Tongue (Rat) Lysate at 40 ug Primary: Anti-MYF5 (bs-6936R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 28 kD Observed band size: 28 kD



Paraformaldehyde-fixed, paraffin embedded (Rat tongue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MYF5) Polyclonal Antibody, Unconjugated (bs-6936R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MYF5) Polyclonal Antibody, Unconjugated (bs-6936R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

- SELECTED CITATIONS -

- [IF=8.3] Cai Bolin. et al. MYH1G-AS is a chromatin-associated lncRNA that regulates skeletal muscle development in chicken. CELL MOL BIOL LETT. 2024 Dec;29(1):1-25 WB ;Chicken. 38177995
- [IF=5.924] Jing Zhang. et al. LncRNA SMARCD3-OT1 Promotes Muscle Hypertrophy and Fast-Twitch Fiber

Transformation via Enhancing SMARCD3X4 Expression. INT J MOL SCI. 2022 Jan;23(9):4510 WB ; Chicken. 35562902

- [IF=3.998] Renli Qi. et al. The intestinal microbiota contributes to the growth and physiological state of muscle tissue in piglets. Sci Rep-Uk. 2021 May;11(1):1-14 WB ; Pig. 34045661
- [IF=2.9] Wesley K R. et al. Effects of implant status and breed type on feedlot performance, carcass characteristics, sera metabolites, and immunohistochemical responses in finishing steers. J ANIM SCI. 2025 Jan;: IHC ;Steers. 39760134