

bs-11042R**[Primary Antibody]****Bestrophin 3 Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

<p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>GeneID: 144453</p> <p>Target: Bestrophin 3</p> <p>Immunogen: KLH conjugated synthetic peptide derived from human Bestrophin 3: 45-92/668.</p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: Bestrophin-3 is a 668 amino acid member of the Bestrophin family of proteins. Members of the Bestrophin family are transmembrane proteins that contain a high percentage of aromatic residues, a conserved RFP (arg-phe-pro) motif and they function as anion channels. Expressed predominantly in skeletal muscle but also found in spinal cord, brain, testis, thymus, retina and bone marrow, Bestrophin-3 forms calcium-sensitive chloride channels. In addition, Bestrophin-3 contains an AI (auto-inhibitory) domain that is capable of regulating the anion channel activity. Due to alternative splicing events, three isoforms exist for Bestrophin-3.</p>	<p>Isotype: IgG</p> <p>SWISS: Q8N1M1</p>	<p>Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000)</p> <p>Reactivity: (predicted: Human, Mouse, Rat, Pig, Dog, Horse)</p> <p>Predicted MW.: 76 kDa</p> <p>Subcellular Location: Cell membrane</p>
--	--	--