

**bsm-60929R****[ Primary Antibody ]****BioSS**  
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**MMP7 Recombinant Rabbit mAb****— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> WB (1:500-2000)
<b>Clonality:</b> Recombinant	<b>CloneNo.:</b> 9F8	<b>Reactivity:</b> Human
<b>GeneID:</b> 4316	<b>SWISS:</b> P09237	
<b>Target:</b> MMP7		
<b>Purification:</b> affinity purified by Protein A		<b>Predicted MW.:</b> 30 kDa
<b>Concentration:</b> 1mg/ml		<b>Subcellular Location:</b> Secreted ,Extracellular
<b>Storage:</b> PBS, Glycerol, BSA Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded by this gene degrades proteoglycans, fibronectin, elastin and casein and differs from most MMP family members in that it lacks a conserved C-terminal protein domain. The enzyme is involved in wound healing, and studies in mice suggest that it regulates the activity of defensins in intestinal mucosa. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3. [provided by RefSeq, Jul 2008]		

**— VALIDATION IMAGES —**

25 ug total protein per lane of various lysates (see on figure) probed with MMP7 monoclonal antibody, unconjugated (bsm-60929R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.